

# Criteria of Test against Absence of Trend in Dispersion Characteristics\*

I.V. Veretel'nikova

Department of Theoretical and Applied Informatics  
Novosibirsk State Technical University  
Novosibirsk, Russia  
ira-veterok@mail.ru

B.Yu. Lemeshko

Department of Theoretical and Applied Informatics  
Novosibirsk State Technical University  
Novosibirsk, Russia  
Lemeshko@ami.nstu.ru

**Abstract** — Distributions of various parametric and nonparametric tests are studied using methods of statistical modeling. Such tests are designed to test hypotheses for randomness or absence of a trend in dispersion characteristics. Respective studies are carried out for different sample sizes with the tested hypothesis being true. Procedure of interactive simulation of distributions of the test statistics is proposed and implemented. Such procedure allows valid usage of corresponding tests in case of violation of standard assumptions. The results of comparative analysis of the power of the tests against competing hypotheses with different models of linear, periodic, and mixed trend in the distribution characteristics are shown. Preferences for using different criteria are given.

**Keywords** — trend, hypothesis of randomness, statistical modeling, power of a test.

## I. INTRODUCTION

A variety of parametric and nonparametric tests has been proposed at different times to test the hypothesis for randomness or absence of a trend in the mathematical expectation and in the dispersion characteristics. However, available sources do not allow us to judge the benefits of a particular test and do not contain any distinct recommendations on the area of application and prerequisites providing correctness of statistical conclusions when using the tests under consideration. As a rule, assumption of normal distribution law of noise is the main prerequisite for ensuring the correct application of parametric tests, but it is not always realized in practice. The usage of nonparametric tests is based on asymptotic distribution of statistics of such tests. For limited sample number distribution of the statistics of parametric and non-parametric tests may differ significantly from the corresponding limit distributions of statistics used for testing the hypothesis. In case of nonparametric tests the problem gets even worse because of apparent discreteness of the statistic. In such situations, the usage of the limiting (asymptotic) distribution of the statistics instead of the true distribution of such statistics may lead to wrong conclusion.

In this paper methods of statistical modeling are used to investigate the distributions and powers of statistical tests designed to check the hypothesis against the absence of a trend in dispersion characteristics (in variance) of observed

random number sequence (measured values).

## II. OVERVIEW OF THE TESTED STATISTICAL HYPOTHESES

When checking the absence of a trend in the dispersion characteristics the task is formulated as follows. It is assumed that time series  $x_1, x_2, \dots, x_n$  of mutually independent random variables is observed. The hypothesis  $H_0: \sigma_i = \sigma$ ,  $i = 1, 2, \dots, n$ , is tested that all sample values belong to the same population with mean  $\sigma$ , against a competitive hypothesis about the presence of a trend  $H_1: |\sigma_{i+1} - \sigma_i| > 0$ ,  $i = 1, 2, \dots, n-1$ .

When checking the absence of a variance shift (in dispersion characteristics) it is assumed that the observed sequence of measured values  $x_1, x_2, \dots, x_n$  has one mean  $\mu$ . The hypothesis  $H_0: \sigma_1^2 = \dots = \sigma_n^2 = \sigma_0^2$  ( $\sigma_0^2$  being unknown) is tested against competitive hypothesis

$$H_1: \sigma_1^2 = \sigma_2^2 = \dots = \sigma_k^2 = \sigma_0^2; \\ \sigma_{k+1}^2 = \sigma_{k+2}^2 = \dots = \sigma_n^2 = \sigma_0^2 + \delta; \quad (\delta > 0)$$

for variance value changes in some unknown point, i.e.  $k$  is unknown ( $1 \leq k \leq n-1$ ).

Analysis of the power of the tests was carried out for the situation when observed random variables belong to the normal law. The tested hypothesis  $H_0$  corresponds to true assumption on independency of observed random values (in other words, the absence of trend). Various situations corresponding to the presence of a trend in the variance were used as competing hypotheses.

During analysis of test powers for the tests against variance change in an unknown point hypotheses close to the  $H_0$  (in case of normal distribution of random variables) were treated as competitive, when at some point the standard deviation was increased by 5, 10, 15%:

$$H_1: \sigma_1^2 = \dots = \sigma_k^2 = 1; \quad \sigma_{k+1}^2 = \dots = \sigma_n^2 = 1.1025, \\ H_2: \sigma_1^2 = \dots = \sigma_k^2 = 1; \quad \sigma_{k+1}^2 = \dots = \sigma_n^2 = 1.21,$$

This work is supported by the Russian Ministry of Education and Science (project 2.541.2014K).

$H_3$ :  $\sigma_1^2 = \dots = \sigma_k^2 = 1$ ;  $\sigma_{k+1}^2 = \dots = \sigma_n^2 = 1.3225$ ,  
where  $k = n/2$ .

One competitive hypothesis was considered as more distant:

$$H_4$$
:  $\sigma_1^2 = \dots = \sigma_k^2 = 1$ ;  $\sigma_{k+1}^2 = \dots = \sigma_n^2 = 4$ .

As the time series that correspond to hypothesis  $H_1$  are visually similar to the time series corresponding to hypotheses  $H_2, H_3$ , Fig. 1 shows only time series for  $H_1$ .

The presence of a linear trend in the dispersion characteristics of the observed series of random variables (change in scale parameter) in the interval  $t \in [0, 1]$  can be simulated according to

$$x_i = \xi_i(1 + ct_i), \quad (1)$$

where  $c \in (-1, \infty)$ ,  $t_i = (i-1)\Delta t$ ,  $\Delta t = 1/n$ . True tested hypothesis  $H_0$  corresponds to parameter value  $c = 0$ .

In case of a periodic trend in the characteristics of dispersion, random values can be simulated, for example, in accordance with the following formula:

$$x_i = \xi_i(1 + d \cdot \sin(2k\pi t_i)) \quad (2)$$

for  $|d| < 1$ . In case of a combined trend it can be simulated according to

$$x_i = \xi_i(1 + ct_i + d \sin(2k\pi t_i)), \quad (3)$$

for  $|d| < 1$ , if  $c \geq 0$ , and for  $|d| < 1 + c$ , if  $c \in (-1, 0)$ . The absence of a periodic component of the trend corresponds to the parameter value  $d = 0$ , and the absence of a linear component corresponds to  $c = 0$ .




Fig. 1. Trend corresponding to  $H_0, H_1, H_4$  hypotheses

During the analysis of power with respect to linear, periodic, and combined trend in the dispersion characteristics (in variance) of a random variable the following competitive hypotheses were considered:

$$H_5$$
:  $x_i = \xi_i(1 + ct_i)$ ,  $c = 1$ ;

$$H_6$$
:  $x_i = \xi_i(1 + d \cdot \sin(2k\pi t_i))$ ,  $d = 0.8$ ,  $k = 2$ ;

$$H_7$$
:  $x_i = \xi_i(1 + ct_i + d \sin(2k\pi t_i))$ ,  $c = 1$ ,  $d = 0.8$ ,  $k = 2$ ;

Representation of corresponding processes is shown in Figures 2 – 4.




Fig. 2. Linear trend in the dispersion characteristics for  $H_5$




Fig. 3. Periodic trend in the dispersion characteristics for  $H_6$




Fig. 4. Combined trend in the dispersion characteristics for  $H_7$

The following tests were compared by powers against competing hypotheses  $H_1 \div H_7$ .

### III. TESTS AGAINST TREND IN DISPERSION CHARACTERISTICS

#### A. Foster-Stuart test

This nonparametric test can be used to test hypotheses of absence of a trend in the mean values or in the variances (dispersion characteristics) depending on the used statistics type. Test against trend in distribution characteristics is given by [1]:

$$S = \sum_{i=2}^n S_i, \quad (4)$$

where  $S_i = u_i + l_i$ ;

$u_i = 1$ , if  $x_i > x_{i-1}, x_{i-2}, \dots, x_1$ , otherwise  $u_i = 0$ ;

$l_i = 1$ , if  $x_i < x_{i-1}, x_{i-2}, \dots, x_1$ , otherwise  $l_i = 0$ .

It is clear that  $0 \leq S \leq n-1$ .

In the absence of a trend the normalized statistics

$$\tilde{t} = \frac{S - \mu}{\hat{\sigma}_S}, \quad (5)$$

where

$$\mu = 2 \sum_{i=2}^n \frac{1}{i}, \quad \hat{\sigma}_S = \sqrt{\mu - 4 \sum_{i=2}^n \frac{1}{i^2}} \approx \sqrt{2 \ln n - 3,4253},$$

are approximately described by Student's distribution with  $\nu = n$  degrees of freedom. The hypothesis of absence of a trend is rejected at large modulus values of statistics (5).

Actually, the area of discrete values is the range of definition of  $\tilde{t}$  statistics. The analysis of statistics distributions shows that even with relatively large sample sizes (around  $n=100, 200$ ) the discrete distributions of test statistics are significantly different from the Student distribution with  $n$  degrees of freedom [2].

#### B. Cox-Stuart test

Cox-Stuart test against the hypothesis of absence of a trend in variance (in dispersion characteristics) is designed as follows.

Initial sample  $x_1, \dots, x_n$  is divided into  $[n/k]$  subsamples with  $k$  number of elements  $x_1, \dots, x_k$ ;  $x_{k+1}, \dots, x_{2k}$ ;  $x_{2k+1}, \dots, x_{3k}$ ;  $\dots$ ;  $x_{n-k+1}, \dots, x_n$  (if  $n$  is not divided by  $k$ , then the required number of measurements in the center is dropped out). For every  $i$  th subsample the range  $w_i$  is found ( $1 \leq i \leq r$ ,  $r = [n/k]$ ). Next, the resulting sequence of ranges is tested against the trend in the mean values using the test with statistics

$$S_1^* = \frac{S_1 - E[S_1]}{\sqrt{D[S_1]}}, \quad (7)$$

where

$$S_1 = \sum_{i=1}^{[n/2]} (n-2i+1)h_{i,n-i+1},$$

$$E[S_1] = \frac{n^2}{8}, \quad D[S_1] = \frac{n(n^2-1)}{24},$$

where  $h_{i,j} = 1$ , if  $x_i > x_j$ , and  $h_{i,j} = 0$ , if  $x_i \leq x_j$  ( $i < j$ ). If the hypothesis for the absence of a trend is true, distribution (7) can be approximately described by standard normal law.

It is recommended to choose the value of  $k$  in [3] according to the following correlations:

$$n \geq 90 \rightarrow k = 5; \quad 64 \leq n < 90 \rightarrow k = 4;$$

$$48 \leq n < 64 \rightarrow k = 3; \quad n < 48 \rightarrow k = 2.$$

The discreteness of the  $S_1^*$  statistics distribution upon detection of a trend in the variance is significantly higher than the discreteness of the Cox-Stuart statistics distribution for trend in mean. This is natural because the analyzed range sample contains only  $[n/k]$  number of elements. When using the Cox-Stuart test for detection of a trend in the dispersion, the difference of statistics discrete distribution from the standard normal law can almost be neglected only for  $n > 170$  [4].

#### C. Hsu test against variance shift and shift point detecting

Under this test the rejection of the hypothesis of randomness (for absence of a trend) can show the discovery of a *variance shift*. Hsu test statistics are given by [5]

$$H = \frac{\sum_{i=1}^n (i-1)(x_i - m_x)^2}{(n-1) \sum_{i=1}^n (x_i - m_x)^2}, \quad 0 \leq H \leq 1, \quad (8)$$

where  $m_x$  is median of variation series. Under the assumption that the mathematical expectation of a sequence of random variables has the same value, the hypothesis of a constant variances is tested. As a competitive hypothesis, the change in the dispersion of observed values at some (unknown) time (starting from some element of the sample) can be considered. The test is two-sided: the tested hypothesis of absence of a variance shift is rejected for small and large values of the statistics (8).

Usually the test is used in a normalized form

$$H^* = \frac{H - 1/2}{\sqrt{D[H]}}, \quad (9)$$

$$\text{where } D[H] = \frac{n+1}{6(n-1)(n+2)}.$$

Under the validity of the hypothesis of the absence of variance changes in the asymptotics, such statistics (9) obeys the standard normal law.

The modeling results [6] show that for  $n > 30$  statistics distribution agrees well with the standard normal law. Statistics distribution (9) strongly depends on the law of distribution to which random variables belong. The greatest deviation from the standard normal law is observed in the case when random variables belong to the laws with heavy tails. Asymmetry of the law significantly affects the statistics distribution.

A test allowing to determine the change point of the variance (in the case when observations belong to the normal law) is proposed in [5]. Statistics of this test are presented as follows. Let for  $k = 1, 2, \dots, n-1$

$$w_k = \sum_{i=1}^k (x_i - m_x)^2, \quad W_k = \frac{w_n - w_k}{w_k} \frac{k}{n-k},$$

where  $k$  corresponds to the required variance change point. If  $x_i$  belongs to normal law, then values of  $W_k$ ,  $k = 1, 2, \dots, n-1$ , belong to corresponding  $F_{n-k,k}(W)$  Fisher distributions with  $n-k$  and  $k$  degrees of freedom.

Next, based on the corresponding distribution functions, we find  $\gamma_k = F_{n-k,k}(W_k)$ , where  $\gamma_k$  must obey to uniform law under the absence of variance shift.

G-test statistics are given by

$$G = \frac{1}{n-1} \sum_{k=1}^{n-1} \gamma_k, \quad 0 \leq G \leq 1. \quad (10)$$

The hypothesis about absence of variance changes is rejected with significance level  $\alpha$ , if  $G < G_{\alpha/2}$  or  $G > G_{1-\alpha/2}$ . In this case value  $k$  corresponding to the maximum value  $|\gamma_k - 1/2|$ , evaluates the desired change point of the variance value in observed series. For  $x_1 = m_x$  value  $w_1 = 0$ , thus  $W_1 = \infty$  and  $\gamma_1 = 1$ . The type of limit distribution of the statistics (10) is not given in the original material, only percentage points are given.

Based on the results of the statistical modeling we have shown that a good model of the limit distribution of the statistics (10) is a beta distribution of the 1st kind with the density of

$$f(x) = \frac{1}{\theta_2 B(\theta_0, \theta_1)} \left( \frac{x - \theta_3}{\theta_2} \right)^{\theta_0 - 1} \left( 1 - \frac{x - \theta_3}{\theta_2} \right)^{\theta_1 - 1}$$

and parameter values  $\theta_0 = 2.7663$ ,  $\theta_1 = 2.7663$ ,  $\theta_2 = 1$ ,  $\theta_3 = 0$  (see Fig. 5). Based on this law we can find percentage points  $G_{\alpha/2}$  and  $G_{1-\alpha/2}$  or  $p$ -values.




Fig. 5. Convergence to the a beta distribution of the 1st kind distribution of G-test statistics

#### D. Klotz and Savage rank tests against variance shifts

Rank tests for detecting the change of the scale parameter (dispersion characteristic) in the unknown point are based on the usage of a family of rank statistics in form [7]

$$S_R = \sum_{i=1}^n i a_n(R_i), \quad (11)$$

where  $R_i$  are ranks of sampled values in an ordered series of measurements.

Tests differ by the used scores  $a_n$ . Their type determines the name of the test. The following scores are commonly used:

- Klotz scores  $a_{1n}(i) = U_{i/(n+1)}^2$ , where  $U_\gamma$  – is a  $\gamma$ -quantile of standard normal law;
- Savage scores  $a_{2n}(i) = \sum_{j=1}^i \frac{1}{n-j+1}$ .

If the tested hypothesis  $H_0$  is true, then tests with statistics

$$S_{R,j} = \sum_{i=1}^n i a_{jn}(R_i), \quad j = 1, 2$$

are symmetric with respect to  $E[S_{R,j}] = \frac{n+1}{2} \sum_{i=1}^n a_{jn}(i)$ .

Usually normalized tests with the following statistics are used

$$S_{R,j}^* = \frac{S_{R,j} - E[S_{R,j}]}{\sqrt{D[S_{R,j}]}} \quad (12)$$

where

$$E[S_{R,1}] = \frac{n+1}{2} \sum_{i=1}^n U_{i/(n+1)}^2, \quad E[S_{R,2}] = \frac{n(n+1)}{2};$$

$$D[S_{R,1}] = \frac{n(n+1)}{12} \sum_{i=1}^n U_{i/(n+1)}^4 - \frac{1}{3n+3} [E[S_{R,1}]]^2$$

$$D[S_{R,2}] = \frac{n(n+1)}{12} \left( n - \sum_{j=1}^n \frac{1}{j} \right).$$

Statistics (12) are approximately obeying the standard law. The convergence of the statistics distributions to the standard law was studied in [6-7].

Statistical modeling research of the distribution of statistics with Klotz scores had shown that for  $n > 20$  distribution is well-approximated by standard normal law. Distribution of the test statistics with Savage scores is also matching well with standard normal law, but only for  $n > 30$ .

#### IV. ANALYSIS OF TEST POWERS

In the course of work statistical simulation methods (for probabilities of errors of the first kind  $\alpha = 0.15, 0.1, 0.05, 0.01$ ) provided estimations of the capacity of the investigated criteria with respect to the competitive hypotheses  $H_1, H_2, H_3$  and  $H_4$  (corresponding to the shift of the dispersion value). The simulation was carried out with the accuracy of  $\pm 10^{-3}$ . Test powers of competing hypotheses  $H_5, H_6, H_7$  were studied. Such hypotheses correspond to linear or nonlinear trend presence in the dispersion characteristics of analyzed processes.

TABLE I. COMPARATIVE ANALYSIS OF POWERS OF ALL RANDOMNESS TESTS AND TESTS AGAINST TREND ABSENCE IN VARIANCE ( $n=100, \alpha = 0.1$ )

№	Against $H_1$	$1 - \beta$	Against $H_2$	$1 - \beta$
1	Hsu H	0.156	Hsu H	0.304
2	Klotz	0.151	Klotz	0.287
3	Hsu G	0.147	Hsu G	0.269
4	Cox-Stuart	0.123	Cox-Stuart	0.188
5	Savage	0.110	Foster-Stuart	0.130
6	Foster-Stuart	0.106	Savage	0.129

№	Against $H_3$	$1 - \beta$	Against $H_4$	$1 - \beta$
1	Hsu H	0.500	Hsu H	1
2	Klotz	0.469	Klotz	1
3	Hsu G	0.430	Cox-Stuart	0.997
4	Cox-Stuart	0.284	Hsu G	0.993
5	Foster-Stuart	0.165	Foster-Stuart	0.625
6	Savage	0.159	Savage	0.610

№	Against $H_5$	$1 - \beta$	Against $H_6$	$1 - \beta$
1	Hsu H	0.836	Hsu H	0.711
2	Hsu G	0.818	Klotz	0.678
3	Klotz	0.807	Hsu G	0.545
4	Cox-Stuart	0.489	Savage	0.196
5	Foster-Stuart	0.346	Cox-Stuart	0.143
6	Savage	0.246	Foster-Stuart	0.048

№	Against $H_7$	$1 - \beta$
1	Hsu H	0.162
2	Savage	0.095
3	Foster-Stuart	0.082
4	Hsu G	0.057
5	Cox-Stuart	0.052
6	Klotz	0.104

Power estimates only with the significance level  $\alpha = 0.1$  and the sample number  $n=100$  are shown in Table 1 for comparative analysis of power. Tests are ordered by

decreasing power  $1 - \beta$ . For similar competitive hypotheses criteria Hsu tests with H and G statistics as well as Klotz test showed the highest power with respect to the analyzed sets of competitive hypotheses. They showed the ability to detect trend in the dispersion characteristics when it has a 10% increase.

Hsu tests with H - and G-statistics and Klotz test can also well distinguish between the null hypothesis and its competitive hypotheses ( $H_5$  and  $H_6$ ) which exhibit the presence of linear or periodic trend in distribution characteristics. At the same time Cox-Stuart, Savage and Foster-Stuart tests can not detect the presence of a periodic trend in the variance reliably (due to relatively low power against similar enough hypothesis  $H_6$ ).

Unfortunately, none of these tests has shown the ability to detect a combined trend in the dispersion corresponding to the studied hypothesis  $H_7$ . The powers with respect to such hypothesis were extremely low.

#### CONCLUSIONS

Thus, methods of statistical modeling have been used to study the statistics distribution of various parametric and nonparametric tests for randomness and the absence of a trend in the dispersion characteristics; within the framework of developing ISW software an interactive study mode of the distributions of the statistics has been implemented for the case of violation of standard assumptions. A comparative analysis of test powers against some competitive hypotheses has been carried out, and results of such analysis can be used to estimate the desirability of application of particular test. Disadvantages of individual criteria have been noted. This research has been supported by the Russian Ministry of Education and Science (project 2.541.2014K)

#### REFERENCES

- [1] Kobzar, A.I. Applied mathematical statistics for engineers and academic researchers/ Kobzar A.I. – M.: Fizmatlit, 2006. – P. 816.
- [2] Veretelnikova, I.V., Lemeshko, B.Yu. Analytical review of tests for randomness and trend absence // Materials of the XII international conference "Actual problems of electronic instrument engineering" (APEIE-2014). V.6, Novosibirsk, 2014. – pp.16-23.
- [3] Cox D.R., Stuart A. Quick sign tests for trend in location and dispersion // Biometrika. 1955. V.42. pp.80–95.
- [4] Veretenikova, I.V., Lemeshko, B.Yu. Application of tests for hypotheses about randomness or trend absence // Cutting-edge technologies, fundamental researches and innovations: сборник статей of the XVII international scientific-practical conference "High Technologies in Industry and Economics" 22-23th May 2014, St. Petersburg, Russia/ Science editors: A.P. Kudinov, M.A. Kudinov. – SPb: Publishing house of the Polytechnic University, 2014. pp. 33-37.
- [5] Hsu D. A. Test for variance shift at an unknown time point / D. A. Hsu // Appl. Statist., 1977. V.26, № 3. P.279–284.
- [6] Lemeshko B.Yu., Komissarova A.S., Tsheglov A.E. Application of tests for trend detection and checking for randomness // Metrologiya. 2010. # 12. – pp. 3-25.
- [7] Lemeshko B.Yu., Komissarova A.S., Tsheglov A.E. Properties and power of tests for trend detection and checking for randomness // Scientific Bulletin of NSTU. – 2012. – # 1(46). – pp. 53-66.